

➤ **Pain Control:** *Tylenol, Ibuprofen, and Hyland's Arnica* are popular pain control medications; these agents may be used at the same time if needed. **Ibuprofen is far superior** to Tylenol in duration, safety, and anti-inflammatory property. *Provide for 2-5 days if needed.*

- **Tylenol** is **NOT effective** post-surgery unless accompanied by opioid. It has **NO ANTI-INFLAMMATORY property** which your baby needs. It is so toxic you are **NOT ALLOWED** to take it for a full day. Many of our patients who have **FAILED** to succeed did so on Tylenol.
- **Ibuprofen** **infant Advil or infant Motrin** every 6 hours; *start soon post-op.*
- **Hyland's Arnica Montana 30X** 1 tablet every 6 hrs; dissolves easy; start soon post-op.

You may choose other infant homeopathic remedies as well but **NOT NUMBING** teething gels.

SUMMARY: **provide pain relief IBUPROFEN for at least 2 days;** unless baby has **kidney problem, dehydrated, or premature.**

WEIGHT	INFANT IBUPROFEN	INFANT TYLENOL
6LBS	0.7 ML	1.3 ML
7LBS	0.8 ML	1.5 ML
8LBS	0.9 ML	1.7 ML
9LBS	1 ML	1.9 ML
10LBS	1.1 ML	2 ML
11LBS	1.2 ML	2.3 ML
12LBS	1.4 ML	2.5 ML
13LBS	1.5 ML	2.7 ML
14LBS	1.6 ML	2.9 ML
15LBS	1.7 ML	3.2 ML
16LBS	1.8 ML	3.4 ML
17LBS	1.9 ML	3.6 ML
18LBS	2 ML	3.8 ML
19LBS	2.1 ML	4 ML
20LBS	2.3 ML	4.2 ML
21LBS	2.4 ML	4.4 ML
22LBS	2.5 ML	4.6 ML
23LBS	2.6 ML	4.8 ML
24LBS	2.7 ML	5 ML